

Journal of Ayurveda Medical Sciences

*Refereed, Indexed, Peer reviewed, Open access, Quarterly
Journal for Rapid Publication of Ayurveda and Other
Traditional Medicine Research*



ISSN: 2456-4990

www.jayumedsci.com

©Journal of Ayurveda Medical Sciences – HRGS' Ayurveda Journal



Pharmacognostic and Phytochemical Analysis of Agnimantha (*Premna corymbosa* Rottl.) Leaf

Rini Sam Susan^{1*}, Harini A², Sunil Kumar KN³, Prakash L Hegde⁴

¹PG Scholar, ²Associate Professor, ⁴Professor, Department of Dravyaguna, Sri Dharmasthala Manjunatheshwara College of Ayurveda, Hassan, Karnataka, 573201. ³SDM Centre for Research in Ayurveda and Allied Sciences, Laxminarayana Nagar, Kuthpadi, Udipi, INDIA 574 118; Current: Research Officer and HOD, Department of Pharmacognosy, Siddha Central Research Institute, Arumbakkam, Chennai, India 600106.

*Correspondence: Email: rinisamsusan89@gmail.com, Mobile: +91-7829077680

ABSTRACT

Introduction: *Agnimantha* (*Premna corymbosa* Rottl.) Family Verbinaceae, a very important plant has been in use since Vedic times. It occurs throughout India, in the plains, as a large perennial shrub, growing to a height of 9 to 10 m. It is one among the combination drugs i.e. *Brhatpanchamoola* (Great 5 root drugs) and of *Dasamoola* (10 root drugs). Its leaves are used as expectorant in asthma, bronchitis, cold, catarrh and fever. In this work an attempt was done to establish pharmacognostic and preliminary phytochemical standards of leaves including HPTLC. **Methods:** Pharmacognostical parameters for the leaves of *P. corymbosa* was performed using parameters like macromorphology, microscopy, physico-chemical constants and phytochemical screening were done using standard methodology. **Results:** The leaves showed presence of tannins and flavanoids. HPTLC densitometric scan showed twelve chemical components at 254nm, nine at 366nm and ten at 620nm. **Conclusion:** The present study will serve as a standard for the identity of *Premna corymbosa* Rottl leaves.

KEYWORDS

Agnimantha, *Brhatpanchamoola*, *Dasamoola*, *Premna corymbosa* Rottl., Standardisation.

Received: 16.10.2016

Accepted: 11.01.2017

DOI: 10.5530/jams.2016.1.14

Agnimantha, *Premna corymbosa* Rottl. (Verbinaceae) is very important plant since Vedic period.¹ It is called as *Agnimathanaha*,^{2,3,4} *Arani*,^{5,6,7} and *Nadeyi*^{4,5,7,8,9} in Sanskrit, *Munja* in Malayalam^{7,10} and *Taggiberu*^{7,10} in Kannada. The twigs of this plant used to be rubbed together to light fire¹ during ceremonial sacrifices, which is evident from the Sanskrit name, *Agnimantha*. It possesses *Kapha Vatahara*¹¹ karma and *Sothahara* (anti-inflammatory) property. In Ayurveda, the usage of *Agnimantha* leaves^{12,13} is mentioned, as stomachic³ and against dyspepsia.³ Leaves are reported to have anti-inflammatory¹⁵, antioxidant¹⁶, hepatoprotective¹⁷, antidiabetic actions¹⁸. The acute and chronic anti-inflammatory activity of leaf ethanolic extract of *Agnimantha* at the doses of 200 and 400mg/Kg body weight showed inhibition in odema formation significantly in acute models, and reduction in the granuloma formation with percentage of 35.17% and 50.38% inhibition in chronic models.¹⁵ Pledging antioxidant action was found from the methanol leaf extract of *P. corymbosa*.¹⁶ Evident hepatoprotective activity and reduction in the liver weight was seen in *P. corymbosa* leaves against carbon tetrachloride (CCl₄) induced hepatic damage.¹⁷ Antidiabetic action of the methanol extract of leaves of *P. corymbosa* was tested by subjecting it with, ethyl acetate and hexane, and the results proves to manage hyperglycemia.¹⁸ Literature on pharmacognostic standards for *P. corymbosa* Rottl. leaf is not available. Considering its wide availability and therapeutic utility, the present work was taken up to establish standardization of *P. corymbosa* Rottl. leaf with the main objective to evaluate pharmacognostic parameters such as macroscopic, microscopy, physicochemical, phytochemical and chromatographic studies of the leaves.

MATERIALS AND METHODS

The leaves of *P. corymbosa* were collected from the herbal garden of Sri Dharmasthala Manjunatheshwara College of Ayurveda and Hospital, Hassan. Authentication was done at the Department of Dravyaguna, Sri Dharmasthala Manjunatheshwara College of Ayurveda and Hospital, Hassan. Authenticated samples were subjected for pharmacognostic, physico-chemical, phytochemical analysis and HPTLC profiling at Department of Pharmacognosy, SDM Centre for Research in Ayurveda and Allied Sciences, Udipi.

Pharmacognostic Study: Macroscopic / organoleptic evaluation of leaves was done for evaluation of external morphology, its shape and size, colour, odour and taste of the drug.¹⁰ A pinch of powder was warmed with drops of chloral hydrate on a microscopic slide and mounted in glycerine. Slides observed under microscope and diagnostic characters were observed and photographed using Zeiss AXIO trinocular microscope attached with Zeiss Axiocamera under bright field light. Magnifications of the figures are indicated by the scale-bars.¹⁰

Physico-Chemical Constants: Physico-chemical constants of leaves were determined to elicit loss on drying, total ash, acid Insoluble ash, water soluble ash, alcohol soluble extractive value and water soluble extractive value.¹⁰

Phytochemical Screening: Phytochemical analysis of leaves were determined to see the presence of alkaloids, steroids, carbohydrate, tannin, flavanoids, saponins, triterpenoids, coumarins, phenols, carboxylic acid, amino acid, resins and quinone.¹⁹

HPTLC Profile: One gram of powdered samples were dissolved in 10 ml of ethanol and kept for cold percolation for 24h and filtered. 3, 6 and 9 µl of the leaf samples of the drug *P. corymbosa* Rottl. were applied on a pre-coated silica gel F254 on aluminium plates to a band width of 7 mm using CAMAG (Muttentz, Switzerland) Linomat 5 TLC applicator. The plate was developed in the solvent system of toluene ethylacetate and formic acid (7.0: 3.0: 0.3) in CAMAG twin trough chamber. The developed plates were visualized in under short and long UV and then derivatised with vanillin sulphuric acid reagent, prior to derivatisation in CAMAG Photo documentation unit. The plate was scanned under UV 254 and 366 nm using CAMAG Scanner 4. R_f, colour of the spots and densitometric scan were recorded.²⁰

RESULTS AND DISCUSSION

The study on the pharmacognostic features of medicinal plants is a process to know its quality, purity and also to check the presence of adulterants and substitutes.

Macroscopical / Organoleptic: Macroscopically the leaves were simple, oblong to obovate in shape, opposite/ whorly arranged; its surface was coriaceous, dark green above and below dull colour, with strong characteristic odour and astringent taste.

Powder microscopy: Leaf powder was rough, light green, strongly aromatic and astringent in taste. In microscopic view, the powder showed the presence of fragment of epidermis with stomata, upper epidermis with palisade, lower epidermis, vein islet, glandular trichome, vasculature in the lamina, thick walled trichome, parenchyma from petiole, thin-walled multicellular trichome, spiral vessels, bundle of fibres and fragment of vasculature (Figure 1).

Physico- chemical evaluation: Determination of loss on drying helps to evaluate the moisture content of a drug. It aids to prevent the decomposition of the drugs either due to chemical change or microbial contamination. From the results obtained for loss on drying, *Agnimanth*a leaves have 9.01% of moisture content.

Determination of ash values is the criterion to judge purity of crude drugs. The residue remaining after incineration is the ash content of the drug. These could be inorganic salts such as carbonates, sulphates, phosphates, silicates etc. naturally occurring in the drug or adhered to it or deliberately added to it in order to adulterate the drug. Since the drugs were collected personally from their natural habitat there was no scope for adulteration. Acid insoluble ash or water soluble ash content is the residue obtained after boiling the total ash either with dilute hydrochloric acid or water which measures the amount of sand and silica matter present in the drug. From the above results the total ash value for leaves of *Agnimanth*a showed 8.01 %, acid insoluble ash was 0.60 %, while water soluble ash content was found to be 3.89 %. Determination of extractive value measures the nature of the chemical constituents present in a crude drug. The results of ethanol soluble and water soluble extractive values of *Agnimanth*a leaves showed 6.11% and 19.18% respectively (Table 1).

Table 1. Physical analysis of leaves of *Agnimanth*a

Parameter	Result
Loss on drying	9.01
Total Ash	8.01
Acid Insoluble Ash	0.60
Water soluble Ash	3.89
Alcohol soluble extractive value	6.11
Water soluble extractive value	19.18


Phyto-chemical evaluation

The aqueous extract of *P. corymbosa* leaves showed the presence of Tannins, Flavanoids, Coumarins and Phenols (Table 2).

Table 2. Preliminary phytochemical screening of leaves of *Agnimanth*a

Class	Result
Alkaloid	-
Steroid	-
Carbohydrate	-
Tannin	+
Flavanoids	+
Saponins	-
Terpenoid	-
Coumarins	+
Phenol	+
Carboxylic acid	-
Amino acids	-
Resins	-
Quinone	-

Figure 1. Powder microscopy of leaves of *Agnimatha*



HPTLC

Retention factors (R_f) values of bands obtained were calculated by exposing the plates to different wavelengths of light. Under short UV, leaves of *P. corymbosa* showed green coloured spots at 0.30, 0.50, 0.59, 0.84 and 0.91. The R_f values in long UV showed red coloured spots at 0.06, 0.27, 0.33, 0.44, 0.55, 0.71, 0.76, 0.82 and 0.91. After derivatisation, when the plate was observed under white light, it showed violet and blue coloured spots for leaves. R_f values in derivatised plate were 0.05, 0.19, 0.27, 0.31, 0.40, 0.44, 0.48, 0.54, 0.59, 0.66, 0.76, 0.84, in that the R_f value 0.84 showed blue coloured spot and the rest was violet coloured (Table 3).

Figure 2. HPTLC of ethanoilc extract of leaves of *Agnimatha*




Figure 3. Densitometric scan of leaves of *Agnimatha*




Table 3. R_f values of leaves of *Agnimatha*

Short UV	Long UV	Post derivatisation
-	-	0.05 (L. violet)
-	0.06 (F. red)	-
-	-	0.19 (L. violet)
-	0.27 (F. red)	0.27 (L. violet)
0.30 (L. green)	-	0.31 (L. violet)
-	0.33 (F. red)	-
-	-	0.40 (L. violet)
-	0.44 (FD. red)	0.44 (L. violet)
-	-	0.48 (L. violet)
0.50 (L. green)	-	-
-	0.55 (FL. red)	0.54 (L. violet)
0.59 (D. green)	-	0.59 (L. violet)
-	-	0.66 (D. violet)
-	0.71 (FD. red)	-
-	0.76 (FD. red)	0.76 (D. violet)
-	0.82 (FD. red)	-
0.84 (L. green)	-	0.84 (L. blue)
0.91 (D. green)	0.91 (FD. red)	-

D – dark; F – fluorescent; L - light

From the results of HPTLC densitometric scan, twelve chemical components at 254nm, nine chemical components at 366nm and ten chemical compounds at 620 nm were found in leaves of *P. corymbosa* (Figure 2 and 3).

CONCLUSION

In the present study, the presence of fragment of epidermis with stomata, glandular trichome, thick walled trichome, spiral vessels, bundle of fibres and fragment of vasculature serve as a microscopic reference standard for leaf identification. The aqueous extract of *P. corymbosa* leaves showed the presence of tannins, flavanoids, coumarins and phenols. The parameters thus obtained from the present pharmacognostic study and HPTLC analysis of the leaves of *Agnimantha* serves as a reference standard for its identity and authenticity.

CONFLICTS OF INTEREST

Nil

REFERENCES

- Sastry JLN. Madanapala Nighantu. 1st ed. Varanasi: Chaukhamba Orientalia; 2010; p.33-35.
- Veel DK, Krishnan R. Tropical Indian Medicinal Plants; Propagation Methods. 1st ed. Bangalore: Polester Prints Pvt. Ltd. 2000; p.285- 6.
- Das PN, Purohit SS, Sharma AK, Kumar T. A Hand Book of Medicinal Plants, A Complete Source Book. Jodhpur: Agrobios (India); 2013; p.421-2.
- Bhandari CR. Vanaushadi chandrodaya. Samskarana; Sapthamam; vim. Sam. 1st part. Varanasi: Chaukhamba Sanskrita Samstana; p.78-9.
- Kirtikar KR, Basu BD. Indian Medicinal Plant. 1st ed.; Vol II. Dehradun: International book distributors; 1995; p.1926-45.
- Kolammal M. Pharmacognosy of Ayurvedic drugs. Series 1; Number 2. Thiruvananthpuram: Pharmacognosy Dept, Ayurveda College; 1978; p.23-4.
- Gupta AK, Tandon N, Sharma M (ed). Review on Indian Medicinal Plants. Vol 7. New Delhi: Indian Council of Medical Research; 2008; p.132-3.
- Database on Medicinal Plants used in Ayurveda. 1sted. Vol II. New Delhi: Central Council for Research in Ayurveda, Department of AYUSH, Ministry of Health & Family Welfare, Government of India; 2005; p.1.
- Pandey G. Dravya-Guna Vijanan. Ist ed. Vol I. Varanasi: Krishnadas Academy; 2001; p.72-6.
- The Ayurvedic Pharmacopoeia of India. Part 1, Vol III, 1sted. New Delhi: Ministry of health and family welfare, Govt. of India; 2001; p.3-4, p.228-30, p.234-7.
- Sharma PV. Dravyaguna Vijnana. Vol I. Varanasi: Chaukhamba Bharati Academy; 2007; p.3.
- Murthy KRS (ed). Astanga Samgraha of Vagbhata. 2nd ed. Vol II. Varanasi: Chaukhamba Orientalia; 1999; p.228-31,481-6.
- Gupta KA, Upadhyaya Y (ed). Ashtanga Hridayam of Vagbhata. 10th ed. Varanasi: Chaukhamba Sanskrit Sansthan; 1987; p.268-9.
- Dey AC. Indian medicinal plants used in Ayurvedic preparation. Dehra Dun: Bishen Singh, Mahendra Pal Singh; 1994; p.1.
- Karthikeyan M, Deepa Karthikeyan. Anti-inflammatory activity of *Premna Corymbosa* (Burm. F.) Rottl & Willd Leaves Extracts in Wistar Rats. Asian Pacific Journal of Tropical Medicine 2011;412-20.
- Radhika S, Senthilkumar R, Arumugam P. Appraisal of In Vitro Antioxidant Prospective of *Premna corymbosa*. International Research Journal of Biological Sciences 2014;3(10):70-5.
- Karthikeyan M, Deepa Karthikeyan. Hepatoprotective Effect of *Premna corymbosa* (Burm.F.) Rottl. & Willd. Leaves Extract on CCl₄ induced Hepatic Damage in Wistar albino Rats. Asian Pacific Journal of Tropical Medicine 2010;3(1):17-20.
- Radhika S, Senthil Kumar R, Sindhu S, Sagadevan E, Arumugam P. Phyto chemical investigation and Evaluation of Antihyperglycemic Potential of *Premna corymbosa*. International Journal of Pharmacy and Pharmaceutical Sciences 2013;5(4):352-6.

19. Agrawal SS, Paridhavi M. Herbal Drug Technology. 1st ed. University Press. 2007; p.321-489.
20. Tandon N, Sharma P. Phytochemical reference standards of selected Indian medicinal plants. Vol 2. New Delhi: Medicinal Plant Unit, Indian Council of Medical Research; 2012; p.330.

ABOUT AUTHORS


Dr. Rini Sam Susan currently working as an Ayurvedic practitioner at Arsha Ayurveda Hospital, Pookulam, Thiruvananthapuram, she accomplished her post graduation in Dravyaguna specialty from Sri Dharmasthala Manjunatheshwara College of Ayurveda & Hospital, Hassan, Karnataka and her under graduation from Mannam Ayurveda Co-operative Medical College, Pandalam, Pathanamthitta, Kerala.

Dr. Harini A is a practicing Ayurvedic physician and is presently positioned as Associate professor in Sri Dharmasthala Manjunatheshwara College of Ayurveda & Hospital, Hassan, India. She is coauthor of a book entitled "Textbook of Dravyaguna (Ayurvedic Clinical Pharmacology)"

Dr. KN Sunil Kumar PhD is currently working as Research Officer in Pharmacognosy at Siddha Central Research Institute, Anna Hospital Campus, Arumbakkam, Chennai. Prior to that, he has worked as Senior Research Officer in Pharmacognosy and Phytochemistry at SDM Ayurveda and Allied Sciences, Udipi, India 574118. He obtained Senior Research Fellowship from ICMR, Young Scientist Award, VGST, Govt. Of Karnataka and Dr. PD Sethi award for 5 best HPTLC papers. He is investigating projects on standardization of Ayurvedic formulation from agencies like UGC, VGST, RGUHS and PCIM (AYUSH). He is Author of 69 research papers and 55 monographs on pharmacognosy, phytochemistry and standardization of medical plants/products. He is also serving as Chief editor Journal of Ayurvedic and Herbal medicine and subject editor Pharmacognosy Ayu-An international Quarterly Journal of Research in Ayurveda.

Dr. Prakash L Hegde obtained his PhD from Dr Sarvepalli Radhakrishnan Rajasthan Ayurveda University, Jodhpur in 2010. He has published a book entitled "Textbook of Dravyaguna (Ayurvedic Clinical Pharmacology)". He is a practicing Ayurvedic physician and specializes in Ayurvedic management of Diabetes mellitus. Dr. Hegde is presently serving as Professor in Sri Dharmasthala Manjunatheshwara College of Ayurveda & Hospital, Hassan, India.

GRAPHICAL ABSTRACT



Cite this article as: Rini Sam Susan, Harini A, Sunil Kumar KN, Prakash L Hegde. Pharmacognostic and Phytochemical Analysis of Agnimantha (*Premna corymbosa* Rottl.) Leaf. J Ayu Med Sci 2016;1(1):80-5. DOI:10.5530/jams.2016.1.14

